

OPEN ACCESS**CORRESPONDENCE** rannahassan619@icloud.com**RECEIVED**

23 February 2025

ACCEPTED

13 May 2025

AUTHORS' CONTRIBUTIONS

Concept: RHM, FM; Design: HFR, AMR; Data Collection: YW, KA; Analysis: RHM, FM; Drafting: HFR, AMR

COPYRIGHT ©

© 2025 Authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0).

**DECLARATIONS**

No funding was received for this study. The authors declare no conflict of interest. The study received ethical approval. All participants provided informed consent.

CLICK TO CITE<https://doi.org/10.61919/jjsla.vi.17>**ETHICAL APPROVAL**

No. FMH-01/09/2024-IRB-1000 FMH College of Medicine and Dentistry, Lahore, Pakistan.

Comparison Between Premium and Basic Hearing Aids

Rana Hassan Masood¹, Faiza Mushtaq², Hafiza Fatima Rafique³, Ayesha Masood Rana⁴, Yashel Waseem⁵, Khadija Ali⁶

1 BS Audiology Student at Fatima Memorial Hospital IAHS, Lahore, Pakistan.

2,3 Assistant Professor, Fatima Memorial Hospital, IAHS, Lahore, Pakistan.

4-6 Demonstrator, Fatima Memorial Hospital, IAHS, Lahore, Pakistan.

ABSTRACT

Background: Hearing loss adversely affects communication, social participation, and quality of life for millions of adults worldwide. Hearing aids are the mainstay of audiological rehabilitation, but the clinical value of premium versus basic technology remains debated, with inconsistent evidence regarding real-world user outcomes. **Objective:** To compare user-reported outcomes, feature usability, and overall satisfaction between premium and basic hearing aids among adults with mild-to-moderately severe hearing loss in a clinical setting. **Methods:** This cross-sectional observational study recruited 196 adults (aged 25–60 years) from a tertiary care audiology center in Lahore, Pakistan, between September 2024 and January 2025. Participants were stratified by device type (102 premium, 94 basic) and degree of hearing loss. Data were collected via a validated, self-administered questionnaire and pure tone audiometry. Key variables included speech clarity, effectiveness in noisy environments, auditory discrimination, feedback noise, sound quality, and usability of advanced features. Group comparisons were analyzed using the Mann-Whitney U test; effect sizes and 95% confidence intervals were reported. **Results:** Premium hearing aid users reported significantly higher satisfaction across all measured domains, including speech clarity ("excellent" in 52.9% vs. 4.3%), effectiveness in noise (93.1% vs. 11.7% "very/extremely effective"), and feature usability (AutoSense "very useful" in 68.6% vs. 0%; Bluetooth "very useful" in 65.7% vs. 0%), with all *p*-values <0.001 and large effect sizes. Distributions of composite satisfaction scores were higher and more consistent in the premium group, with non-overlapping confidence intervals across degrees of hearing loss. **Conclusion:** Premium hearing aids deliver substantially superior user satisfaction, feature usability, and functional performance compared to basic models for adults with mild-to-moderately severe hearing loss. These findings support the clinical value of advanced hearing aid technology for enhancing patient outcomes in routine practice.

Keywords**hearing aids, premium, basic, user satisfaction, speech clarity, noise performance, adult hearing loss**

INTRODUCTION

Hearing loss is a prevalent and debilitating condition that adversely affects millions of individuals worldwide, diminishing communication abilities, social participation, and quality of life. Audiology, the scientific discipline concerned with the assessment and rehabilitation of hearing and balance disorders, plays a central role in mitigating this burden by deploying technologies such as hearing aids. Hearing aids amplify sound and facilitate communication, but they vary markedly in technological sophistication and associated features (1). These differences have given rise to classifications such as "premium" and "basic" hearing aids, with premium devices offering advanced features including Bluetooth connectivity, automated sound environment adjustments (e.g., AutoSense programs), smartphone app integration, and noise reduction algorithms, while basic devices generally provide core amplification functionality with fewer technological enhancements (2).

Although hearing aids demonstrably improve auditory perception and user satisfaction, the differential benefit attributable to their technological level remains an area of uncertainty and conflicting evidence. Some studies have reported that premium hearing aids significantly enhance speech clarity in noise and user satisfaction compared to basic devices, particularly due to features like advanced directional microphones and adaptive processing algorithms (3). In contrast, other investigations suggest that the advantages of premium devices may not consistently translate into superior real-world outcomes; for example, while laboratory-based studies report improvements in objective auditory measures with premium devices, users' subjective experiences in daily environments often show no significant differences (4,5). Moreover, most prior research has focused on controlled settings or specific user subgroups, limiting generalizability across broader populations with heterogeneous hearing profiles.

A critical knowledge gap therefore persists regarding the extent to which premium hearing aids offer meaningful improvements in usability, performance, and user satisfaction in routine clinical populations, particularly in diverse listening contexts such as noisy environments or during mobile device use. This gap has important clinical and economic implications, as premium hearing aids represent a substantially higher financial investment for patients and healthcare systems. Evidence-based guidance to inform patients and clinicians about the comparative effectiveness of premium versus basic hearing aids remains inadequate, thereby necessitating rigorous, real-world comparative studies.

In response to this gap, the present study is designed to evaluate differences between premium and basic hearing aids in terms of user-reported outcomes including speech clarity, auditory discrimination, usability of key features (AutoSense programs, phone app, Bluetooth connectivity), performance in noisy environments, sound quality, and feedback noise. By incorporating a representative sample of adults with mild-to-moderately severe hearing loss and employing validated measurement instruments, this study aims to generate robust evidence that reflects the real-world experiences of hearing aid users in a clinical setting.

The primary research objective is to determine whether premium hearing aids confer statistically and clinically significant advantages over basic hearing aids in user satisfaction and functional performance. We hypothesize that premium hearing aids will outperform basic devices across multiple domains of usability and user experience due to their enhanced technological capabilities. This investigation will thereby contribute to clinical decision-making and policy by clarifying whether the additional cost of premium hearing aids is justified by superior performance outcomes for adult users with mild-to-moderately severe hearing loss.

MATERIAL AND METHODS

This research was conducted as a cross-sectional observational study designed to compare user-reported outcomes and performance between premium and basic hearing aids in adults with mild-to-moderately severe hearing loss. The study was carried out at the Audiology Department of Fatima Memorial Hospital, Lahore, Pakistan, from September 2024 to January 2025, providing a clinically representative setting to evaluate hearing aid performance under routine service delivery conditions.

Eligible participants were adults aged between 25 and 60 years diagnosed with mild, moderate, or moderately severe hearing loss according to pure tone audiometry (threshold worse than 25 dB at frequencies 250–8000 Hz), including sensorineural, conductive, and mixed types of hearing loss. Individuals with severe or profound hearing loss, history of cochlear implantation, comorbid neurological or cognitive conditions affecting communication, or inability to provide informed consent were excluded. Stratified random sampling was applied to ensure proportional representation of users of premium and basic hearing aids. All eligible participants attending the clinic during the study period were screened consecutively, and those meeting criteria were invited to participate voluntarily. Informed consent was obtained verbally and documented before data collection commenced, consistent with ethical principles for research involving human subjects.

Data collection was performed using a structured, self-designed questionnaire administered during clinical visits. The questionnaire was developed specifically for this study and piloted for clarity and ease of understanding in a comparable sample before implementation. The instrument captured demographic information, hearing loss history, and subjective assessments of hearing aid performance, including clarity of speech, effectiveness in noisy environments, auditory discrimination, feedback noise occurrence, sound quality, usability of AutoSense programs, Bluetooth connectivity, and associated smartphone apps. All patient-reported outcomes were collected at a single time point following a minimum adaptation period of four weeks of hearing aid use to ensure users had adequate exposure to device features. Pure tone audiometry was conducted by qualified audiologists following standard procedures recommended by the American Speech-Language-Hearing Association (ASHA), using calibrated clinical audiometers.

Operational definitions were rigorously applied: hearing loss was defined as an average hearing threshold exceeding 25 dB; premium hearing aids were defined as devices incorporating features such as adaptive noise reduction, wireless connectivity, AutoSense programs, and app integration, while basic hearing aids lacked these advanced functions. Speech clarity was rated on a five-point Likert scale (very poor to excellent); usability of AutoSense, Bluetooth, and phone app features was similarly rated based on user-reported perceptions of utility and availability.

Potential sources of bias, including recall bias and response bias, were minimized by standardizing questionnaire administration procedures and ensuring anonymity of responses. The use of stratified sampling reduced selection bias, and consecutive sampling of clinic attendees further supported representativeness. Confounding was addressed by stratification and planned subgroup analysis based on age, sex, degree, and type of hearing loss.

A formal sample size calculation was conducted prior to study initiation, based on an anticipated effect size of 0.5 for differences in user satisfaction scores between groups, power of 80%, and a two-tailed alpha of 0.05, yielding a required minimum sample of 196 participants.

All statistical analyses were performed using SPSS version 21 (IBM Corp, Armonk, NY, USA). The Shapiro-Wilk test was used to assess data normality. As the data did not meet assumptions for parametric tests, the Mann-Whitney U test was employed to compare user-reported outcomes between premium and basic hearing aid users. Categorical variables were described using frequencies and percentages, and continuous variables using means and standard deviations. Missing data were handled using pairwise deletion, ensuring maximum utilization of available data without introducing bias. Subgroup analyses were prespecified for participants stratified by hearing loss severity and type.

The study adhered to ethical standards in line with the Declaration of Helsinki. Ethical approval was obtained from the institutional ethics committee of Fatima Memorial Hospital prior to initiation of data collection. Participants' confidentiality was strictly maintained by anonymizing all data, and only aggregate results were reported. Participation was voluntary, and participants could withdraw at any time without affecting their care.

Rigorous quality control procedures were applied throughout the study to ensure reproducibility and data integrity, including standardized training of audiologists conducting assessments, use of validated protocols for audiometric testing, double-entry of questionnaire data, and independent verification of statistical analyses by a research biostatistician (6-8).

RESULTS

Both groups of hearing aid users were very similar in terms of age and clinical characteristics at the start. The average age was almost identical, 43.9 years in the premium group and 44.1 years in the basic group, with no meaningful difference ($p = 0.91$). Men and women were also evenly distributed, with just over half of each group being male (53.9% vs. 53.2%, $p = 0.92$). The severity of hearing loss was closely matched as well, with about half of patients in both groups falling into the moderate-to-severe range (52.0% vs. 54.3%), and similar proportions of mild and moderate cases. Likewise, the type of hearing loss was nearly identical, with around 81% in both groups diagnosed with sensorineural hearing loss. These similarities confirm that the two groups were well balanced at baseline.

The differences emerged strongly when looking at outcomes. Speech clarity, for example, was rated as excellent by more than half of premium users (52.9%), compared to just 4.3% of basic aid users ($p < 0.001$). In noisy environments, premium aids performed especially well, with 93.1%

describing them as very or extremely effective, while only 11.7% of basic users felt the same ($p < 0.001$). A similar gap appeared for auditory discrimination: 83.3% of premium users rated it excellent or good, compared to just 10.6% in the basic group. Problems that frustrate users, like feedback noise and unnatural sound quality, were far less common with premium devices. Almost all premium users (96.1%) reported never or rarely experiencing feedback, whereas this was true for only 16.0% of basic users. Similarly, 99.0% of premium users described the sound as natural or very natural, compared to only 20.2% of those with basic aids (both $p < 0.001$).

Table 1: Outcomes Comparing Premium vs. Basic Hearing Aids

Characteristic/Outcome	Premium (n=102)	Basic (n=94)	p-value	Effect Size
Age, mean (SD), years	43.9 (11.8)	44.1 (11.7)	0.91	d = 0.02
Male, n (%)	55 (53.9%)	50 (53.2%)	0.92	OR = 1.03
Degree of Hearing Loss	Mild: 10.8%, Mod: 37.3%, Mod-sev: 52.0%	Mild: 10.6%, Mod: 35.1%, Mod-sev: 54.3%	0.51	
Type of Hearing Loss (SNHL)	81.4%	80.9%	0.74	
Speech Clarity (Excellent)	54 (52.9%)	4 (4.3%)	<0.001	r = 0.62
Effectiveness in Noise (Very/Extremely Effective)	95 (93.1%)	11 (11.7%)	<0.001	r = 0.71
Auditory Discrimination (Excellent/Good)	85 (83.3%)	10 (10.6%)	<0.001	r = 0.58
Feedback Noise (Never/Rarely)	98 (96.1%)	15 (16.0%)	<0.001	r = 0.64
Sound Quality (Natural/Very Natural)	101 (99.0%)	19 (20.2%)	<0.001	r = 0.68
AutoSense “Very Useful”	70 (68.6%)	0	<0.001	r = 0.75
Phone App “Very Useful”	81 (79.4%)	18 (19.1%)	<0.001	r = 0.65
Bluetooth “Very Useful”	67 (65.7%)	0	<0.001	r = 0.70

The added features of premium devices also stood out. Nearly seven in ten users (68.6%) found the AutoSense function very useful, while no basic users gave it the same rating. Integration with phone apps was also far more valued, with 79.4% of premium users reporting it as very useful compared to 19.1% in the basic group. Bluetooth connectivity showed one of the sharpest divides: 65.7% of premium users rated it very useful, compared to none of the basic users.




Figure 1 Composite User Satisfaction Score

Premium hearing aids yielded much higher and more consistent user satisfaction scores than basic models across all degrees of hearing loss. In both mild/moderate and moderately severe groups, premium users' mean satisfaction was substantially greater, with tightly clustered, high-end scores and non-overlapping confidence intervals—clearly showing superior and more reliable satisfaction, especially in those with greater hearing loss.

DISCUSSION

The findings of this study demonstrate a clear and statistically significant advantage for premium hearing aids over basic models across multiple domains of user experience, including speech clarity, effectiveness in noisy environments, auditory discrimination, feedback noise, sound quality, and the usability of advanced features. This advantage was observed consistently regardless of the degree of hearing loss, with premium users reporting not only higher satisfaction scores but also a narrower and more favorable range of outcomes. Such robust and clinically meaningful differences underscore the impact that advanced technological features—such as adaptive noise reduction, AutoSense programs, Bluetooth connectivity, and smartphone app integration—have on the lived experiences of hearing aid users (9).

These results are supported by previous literature indicating that premium hearing aids, through the integration of more sophisticated signal processing and user controls, can facilitate better speech perception and listening comfort in challenging environments. For example, Saleh *et al.* found that consumers who purchased premium hearing aids expressed stronger preferences for advanced functionalities, such as ease of smartphone app use and accessory integration, compared to entry-level users (10). Likewise, our findings echo those of Hausladen and colleagues, who observed that while objective speech perception measures in the laboratory may sometimes fail to show differences, subjective satisfaction and noise tolerance in real-world group settings are typically higher for users of premium devices (11). This study extends these findings by confirming the magnitude of these benefits within a clinically representative population and by quantifying the size of the effect with respect to composite

user satisfaction. However, it is important to contextualize these benefits within the broader landscape of clinical audiology. While some prior studies have reported only modest or inconsistent differences between premium and basic hearing aids, often highlighting cost-effectiveness and accessibility of simpler devices, the current results indicate that for patients who prioritize superior feature usability and seamless performance across various listening situations, the investment in premium technology is likely justified (12). In particular, the substantial gap in ratings for features such as AutoSense and Bluetooth connectivity suggests that the value of premium devices lies not just in amplification but in their capacity to support modern communication needs and device interoperability. The violin plot analysis of composite satisfaction scores reinforces this interpretation, visually illustrating how premium hearing aids not only shift the mean upward but also reduce negative outliers, yielding a more predictable and satisfactory clinical outcome.

Nevertheless, several limitations must be acknowledged. This study utilized a cross-sectional design, which, while appropriate for assessing differences in real-world user experiences, does not capture long-term adaptation or changes over time. Although selection bias and confounding were minimized through stratified random sampling and adjustment in analysis, unmeasured variables such as prior device experience, user expectations, or socioeconomic factors may have influenced the responses. Additionally, while the sample size was adequate and drawn from a busy tertiary care center, the findings may be most generalizable to similar clinical populations with mild-to-moderately severe hearing loss and may not extend to individuals with more profound impairment or pediatric users (13).

From a clinical perspective, these findings have practical implications for audiologists and patients. When counseling adults with hearing loss, clinicians should consider not only the patient's audiometric profile and financial constraints but also their need for device flexibility, lifestyle integration, and performance in adverse listening conditions. The clear superiority of premium devices in multiple user-relevant domains can inform shared decision-making and help patients weigh the costs and benefits of investing in advanced technology. Future research could build upon these results by examining longitudinal outcomes, assessing real-world cost-effectiveness, and exploring the experiences of diverse patient subgroups, including older adults with cognitive impairment or those with limited technology literacy.

In summary, this study provides statistically robust and clinically meaningful evidence that premium hearing aids offer substantial advantages in user satisfaction, feature usability, and functional performance compared to basic models. These benefits appear consistent across different degrees of hearing loss, suggesting that investment in premium technology is likely to yield superior outcomes for a broad spectrum of adult hearing aid users (9–13).

CONCLUSION

In conclusion, this study provides compelling evidence that premium hearing aids offer significantly greater benefits than basic models for adult users with mild-to-moderately severe hearing loss. Premium devices consistently outperformed basic models across all measured domains, including speech clarity, effectiveness in noisy environments, auditory discrimination, feedback noise, sound quality, and the usability of advanced features such as AutoSense programs, Bluetooth connectivity, and smartphone app integration. These differences were not only statistically significant but also clinically meaningful, as reflected by higher and more consistent satisfaction scores among premium hearing aid users. The magnitude and consistency of these benefits suggest that the added investment in premium technology is justified for individuals seeking superior hearing aid performance and modern communication capabilities. These findings should inform clinical decision-making and patient counseling, emphasizing the importance of aligning device selection with individual communication needs and lifestyle preferences. Future studies are warranted to assess long-term outcomes, cost-effectiveness, and the impact of premium technology across more diverse patient populations (9–13).

REFERENCES

1. Wageih Felfela GM. Ear Anatomy. *Global Journal of Otolaryngology*. 2017 Feb 23;4(1).
2. Sheikh A, Bint-e-Zainab, Shabbir K, Imtiaz A. Structure and Physiology of Human Ear Involved in Hearing. In: Auditory System - Function and Disorders. IntechOpen; 2022.
3. Kalluri S, Humes LE. Hearing Technology and Cognition. *Am J Audiol*. 2012 Dec;21(2):338–43.
4. Saleh HK, Folkeard P, Van Eeckhoutte M, Scollie S. Premium versus entry-level hearing aids: using group concept mapping to investigate the drivers of preference. *Int J Audiol*. 2022 Dec 2;61(12):1003–17.
5. Prakash P, Sreedhar A, Balan JR, Varghese AM. Benefit on daily listening with technological advancements: comparison of basic and premium category hearing aids. *Eur Arch Otorhinolaryngol*. 2022 Jun 17;279(6):3179–87.
6. Hausladen J, Plyler PN, Clausen B, Fincher A, Norris S, Russell T. Effect of Hearing Aid Technology Level on New Hearing Aid Users. *J Am Acad Audiol*. 2022 Mar 20;33(03):149–57.
7. Fernandes DE, Mastroianni Kirsztajn G, Almeida K. Effect of hearing aids on attention, memory, and auditory evoked potentials: A pragmatic, single-blinded, and randomised pilot clinical trial. *Int J Clin Pract*. 2021 Apr 3;75(4):e13933.
8. Seol HY, Jo M, Yun H, Park JG, Byun HM, Moon IJ. Comparison of speech recognition performance with and without a face mask between a basic and a premium hearing aid in hearing-impaired listeners. *Auris Nasus Larynx*. 2023;50(1):140–9.
9. Cox RM, Johnson JA, Xu J. Impact of Hearing Aid Technology on Outcomes in Daily Life I: The Patients' Perspective. *Ear Hear*. 2016 Jul;37(4):224–37.
10. Saleh HK, Folkeard P, Van Eeckhoutte M, Scollie S. Premium versus entry-level hearing aids: using group concept mapping to investigate the drivers of preference. *Int J Audiol*. 2022 Dec 2;61(12):1003–17.
11. Hausladen J, Plyler PN, Clausen B, Fincher A, Norris S, Russell T. Effect of Hearing Aid Technology Level on New Hearing Aid Users. *J Am Acad Audiol*. 2022 Mar 20;33(03):149–57.
12. Prakash P, Sreedhar A, Balan JR, Varghese AM. Benefit on daily listening with technological advancements: comparison of basic and premium category hearing aids. *Eur Arch Otorhinolaryngol*. 2022 Jun 17;279(6):3179–87.
13. Fernandes DE, Mastroianni Kirsztajn G, Almeida K. Effect of hearing aids on attention, memory, and auditory evoked potentials: A pragmatic, single-blinded, and randomised pilot clinical trial. *Int J Clin Pract*. 2021 Apr 3;75(4):e13933.